

DEVICE FOR THE RAPID MEASUREMENT OF ENZYMATIC ACTIVITY


The present invention relates to a device for the rapid measurement of an enzymatic activity in a solid feed, comprising (i) a container designed to contain the test sample, (ii) a reagent specific for the enzyme whose activity it is desired to measure, and (iii) a buffer for dissolving the enzyme.

The feed is preferably a solid feed which is
10 not treated prior to the measurement.

Feeds intended for husbandry animals are usually supplemented with enzymes whose role is mainly to improve the digestibility of the feed ration. These enzymes are usually sprayed in liquid form onto the feeds, in particular as described in patent EP 0,789,291. The enzymes can also be added in powder form to the feed.

Two problems thus arise, the first being to check the uniformity of distribution of the enzymes added to the feed, the second being to quickly and easily evaluate the activity of the enzyme(s) added to the feeds. These problems are raised in particular by feed manufacturers and breeders wishing to check the quality of the feeds they want to give to their animals. Until now, the enzymatic activity could be measured in the laboratory, thus entailing constraints in terms of logistics and delays, these constraints

being a real hindrance when an immediate result is needed.

Sub B' 

The present invention satisfies this problem by providing a device for measuring the enzymatic activity of any enzyme-enriched feed intended for animal feed. This device, whose measurement is based on a colorimetric reaction, allows both a qualitative measurement of the enzymatic activity of the test sample and a semi-quantitative measurement of this sample.

Figure 1 represents one embodiment of the invention in the form of a device for measuring enzymatic activity, which is in the form of a column.

The description below can be read with regard to the figure mentioned above.

The device which is the subject of the present invention comprises a container designed to contain the test sample, a reagent specific for the enzyme whose activity it is desired to measure and a buffer for dissolving the said enzyme.

The container of this device can be, without any implied limitation, a column (Figure 1) composed of a graduated narrow bottom part (11) and a wide funnel-shaped top part (12) for introducing various reagents into the column and for mixing them during stirring. The column can also be fitted with a leakproof opening and closure system (13) such as a stopper attached to the body of the column by means of a tab (131).

The container can also consist of a single-use tube (Figure 2).

The container can be made of synthetic material such as a single-use plastic.

5 The container can preferably comprise a cleavable protuberance (14) at its base, allowing the liquid part of its contents to flow out. A constriction (141) retaining the solid morsels of feed is advantageously mounted on the protuberance.

10 Measurement of the enzymatic activity is based on the coloration reaction of the Azo method. The principle of the coloration reaction of the Azo method is based on the enzymatic hydrolysis of a characteristic substrate of an enzyme linked to a
15 chromophore. The reaction produces soluble oligomers which turn the medium blue. The absorbence of the medium can be measured at 590 nm.

 The reagent used in the device is the substrate of the reaction catalysed by the enzyme
20 linked to a chromophore. Thus, the enzymatic hydrolysis reaction releases the chromophoric substrate.

 The device also comprises a buffer for dissolving the enzymes which have been sprayed onto the feed, and for keeping the enzymes at their optimum pH.

25 Mention may be made, by way of example and without any restriction being implied, of the device for demonstrating the activity of xylanases.

To measure the activity of xylanases, the reagent used is "Oat spelt Xylan Remazol Brilliant Blue R" or "Xylazyme AX" (sold by the company Megazyme and consisting of oat or wheat araboxylane linked to a dye).

The buffer used is chosen from acetic acid/sodium acetate; glycine hydrochloride/glycine; aconitic acid/sodium hydroxide; formic acid/sodium formate buffers.

Mention may also be made of the device for demonstrating the activity of β -glucanases, which is also based on the coloration reaction of the Azo method.

Among the substrates which can be used, mention may be made of 1,3: 1,4- β -D-glucan with Remazol Brilliant Blue R and Beta-Glucazyme sold by the company Megazyme and consisting of beta-glucan combined with azurine.

The buffer used is chosen from acetic acid/sodium acetate; glycine hydrochloride/glycine; aconitic acid/sodium hydroxide; formic acid/sodium formate buffers.

To measure the activity of cellulase, the substrate used is in the form of Cellazyme lozenges (sold by the company Megazyme). These lozenges consist of substrates based on cellulose and/or on cellulose and xyloglucans polymerized with an azurine dye.

In one preferred embodiment of the present invention, the reagent is in a solid form.

Advantageously, to facilitate the dissolution of the enzyme, a surfactant can be added to the
5 substrate containing the chromophoric agent. This surfactant is chosen in particular from sodium lauryl sulphate and sodium dodecyl sulphate.

According to a better embodiment of the invention, the measurement is carried out in four
10 steps:

- introduction into the container (1) of
10 ml of sample whose enzymatic activity it
is desired to measure - for a solid sample,
the container should be filled with solid up
15 to the 10 ml graduation mark;
- introduction of the reagent in the form of
a solid bead;
- introduction of the specific buffer up to
the 20 ml graduation mark;
- 20 - after closing the column with the stopper,
the column is shaken vigorously several
times.

An additional step of separating the liquid phase and the solid phase (by centrifugation or
25 filtration) can optionally be added, to recover the liquid phase and to measure the intensity of the coloration by spectrophotometry or simply by comparison with a colour scale.

The appearance of a blue coloration after a reaction time of 4 to 8 hours confirms the presence of active enzymes, the intensity of the coloration being proportional to the activity of the enzymes present in the sample.

Another advantage of the present invention is the ability to carry out a semi-quantitative measurement of the enzymatic activity. The coloured liquid phase in the column can be recovered by cutting off the cleavable protuberance from the column. The intensity of its coloration can then be compared with an OD calibration curve.

In addition to being fast, the measurement method is very simple and the device can be used anywhere without requiring special equipment. For example, a manufacturer or a breeder can carry out a control measurement as soon as the feed has been manufactured.

The present invention will be described more fully with the aid of the examples which follow, which should not be considered as limiting the invention.

Examples

Two series of tests were carried out on Rovabio xylan LC (mixture of xylanase and beta glucanase from *Penicillium funiculosum*) and on Rovabio xylanase TRLC (xylanase from *Trichoderma reesei*) whose xylanase activity is between 350 and 550 uAXC/ml. It is

estimated that the treatment of spraying the liquid composition on the feeds leads to a level of 70 to 110 uAXC/kg of feed.

The buffer used is the acetate buffer for maintaining a pH of 4.7. The spraying can be carried out on the feed in pulverulent form or in granulated form.

Sample	Activity	Observa- (before adjust- ment)	tion at 3 hours	O.D. at 590 nm at 4h30	O.D. at 590 nm at 8h	Observa- tion at 8h
xylanase TRLC on granules	1336	blue: +++		>3.0 >3.0		blue: +++
xylanase TRLC on granules	886.7	blue: +		1.567 2.685		blue: ++
xylanase TRLC on granules	1469.25	blue: +		1.429 2.652		blue: ++
xylanase powder before granula- tion	631.4	no color- ation		0.201 0.666		blue: +

xylanase	1144	blue:	2.309	2.376	blue:
LC on		+++			+++
granules					
xylanase	1386.7	blue: +	1.382	2.484	blue:
LC on					+++
granules					
xylanase	1450.5	blue:	2.872	2.85	blue:
LC on		+++			+++
granules					
xylanase	1330.5	blue: ++	1.233	2.096	blue:
LC on					+++
granules					